



# Calibration and evaluation of the *STICS* soil-crop model for faba bean to explain variability in yield and N<sub>2</sub> fixation



Gatien N. Falconnier<sup>a,e,f,\*</sup>, Etienne-Pascal Journet<sup>a,b</sup>, Laurent Bedoussac<sup>c</sup>, Anthony Vermue<sup>a</sup>, Florent Chlébowski<sup>d</sup>, Nicolas Beaudoin<sup>d</sup>, Eric Justes<sup>a,g,h</sup>

<sup>a</sup> AGIR, Université de Toulouse, INPT, INP-PURPAN, INRA, 31320, Auzeville, France

<sup>b</sup> LIPM, Université de Toulouse, INRA, CNRS, Castanet-Tolosan, France

<sup>c</sup> AGIR, Université de Toulouse, INPT, INP-PURPAN, INRA, ENSFEA, 31320, Auzeville, France

<sup>d</sup> UR INRA 1158 Agro-Impact, site de Laon, Pôle du Griffon, F-02000 Barenton-Bugny, France

<sup>e</sup> CIRAD, UPR AIDA, F-34398 Montpellier, France

<sup>f</sup> AIDA, Univ Montpellier, CIRAD, Montpellier, France

<sup>g</sup> CIRAD, UMR SYSTEM, F-34398 Montpellier, France

<sup>h</sup> SYSTEM, Univ Montpellier, CIHEAM-IAMM, CIRAD, INRA, Montpellier SupAgro, Montpellier, France

## ARTICLE INFO

### Keywords:

Parameter estimation  
Yield variability  
Abiotic stresses  
Grain legume

## ABSTRACT

Faba bean (*Vicia Faba* L.) is the second most widely grown grain legume in Europe after pea (*Pisum Sativum* L.) and presents several agronomic and environmental advantages when inserted in cropping systems (e.g. decreased dependency on synthetic fertilisers and N provision to the subsequent crop). However, yield variability due to several factors including heat and drought impede wide adoption of faba bean by farmers. Soil-crop models provide quantitative information to evaluate these processes and help to design innovative cropping system including legumes. The *STICS* model was chosen for its agro-environmental purpose and its genericity allowing crop rotation simulation and robustness for a wide range of pedoclimatic conditions. However, there is so far for *STICS* no parameterization for faba bean. We calibrated 38 crop related parameters based on literature, direct measurements and sequential estimation using the optimisation tool *OptimiSTICS* and a dataset of winter faba bean grown in two sites in France with contrasting soil conditions over several growing seasons (2002–2015). Data from 22 experimental plots were used for calibration and the remaining independent 13 plots were used for model evaluation. After calibration, the *STICS* model reproduced adequately phenology, Leaf Area Index and dynamic growth of above ground biomass, uptake of mineral N, N<sub>2</sub> fixation and grain yield, with satisfactory model efficiency (0.56 to 0.81) and low relative bias (–7% to 2%). The model adequately reproduced the large observed variation in faba bean grain yield (0.42–4.70 t ha<sup>–1</sup>) and total N<sub>2</sub> fixed at harvest (62–172 kg N ha<sup>–1</sup>) in the contrasted years and soil conditions of this study. Simulations indicated that water stress was the overriding factor driving yield and N<sub>2</sub> fixation variability. Simulation of temporal crop growth and water stresses during grain onset and grain filling allowed a robust and credible agronomic diagnosis of the causes of this variability for faba bean crops not significantly damaged by pests and diseases. Water supply/demand ratio averaged over a period of six days preceding beginning of grain filling explained 78% of the observed grain yield variability while water deficit factor for N<sub>2</sub> fixation averaged over a period of 20 days following the beginning of grain filling explained 83% of the variability of fixed N<sub>2</sub> at harvest. Our work provides a first calibration and evaluation of the *STICS* model for faba bean. It offers the opportunity to quantify the ecosystem services associated with crop rotations including faba bean and the effect of climate change on the performance of such rotations.

\* Corresponding author. Current address: CIRAD, UPR-Agro-Ecology and Sustainable Intensification of Annual Crops, University of Montpellier, Avenue Agropolis, 34398 Montpellier, France; CIRAD, UMR SYSTEM, Montpellier Univ, CIHEAM-IAMM, CIRAD, INRA, Montpellier SupAgro, 2 place Viala, F-34060 Montpellier, France

E-mail address: [gatien.falconnier@cirad.fr](mailto:gatien.falconnier@cirad.fr) (G.N. Falconnier).

<https://doi.org/10.1016/j.eja.2019.01.001>

Received 8 December 2017; Received in revised form 29 November 2018; Accepted 10 January 2019

Available online 25 January 2019

1161-0301/ © 2019 Elsevier B.V. All rights reserved.

## 1. Introduction

Current cropping systems in Europe are characterized by high productivity but rely on simplification of crop rotations and high level of synthetic inputs which has led to soil, water and air pollution and loss of biodiversity (e.g. Stoate et al., 2009). Diversification of current cropping systems with grain legumes can improve their agronomic, economic and environmental performance and provide supporting and regulating ecosystem services. In particular, relying on biological N<sub>2</sub> fixation decreases the dependence on synthetic nitrogen fertilizer and reduces the associated direct and indirect greenhouse gases emissions and risks of nitrate leaching (Plaza-Bonilla et al., 2017; Reckling et al., 2016). However, legumes often suffer from water and temperature stress: i) drought during reproductive phase can cause severe yield reduction (Daryanto et al., 2017) and ii) heat can impact negatively seed set and grain filling (Prasad et al., 2017). Even though not always well appreciated, N<sub>2</sub> fixation can also be severely restricted by stresses due to drought, heat, anoxia and soil mineral N content in the root zone (Liu et al., 2011). Furthermore, legumes are exposed to damages from pests, diseases and weeds at different time in their growth cycle (Rubiales et al., 2015). Strong yield variability and lack of reliable agronomic diagnosis are the main reasons mentioned by farmers for non-adoption of legumes in cropping systems (Meynard et al., 2017; Watson et al., 2017). Dynamic process-based soil crop models can assist the understanding of current yield variability by simulating patterns of seasonal water and temperature stress (Affholder et al., 2003; Hayman et al., 2010). They are useful tools to diagnose agronomic constraints to crop production and to explore the impact of current and future climate variability. Therefore, they can assist the design of more resilient cropping systems. Furthermore, these models allow to estimate indicators that are crucial for the assessment of crop rotations but tedious or expensive to quantify in experimental plots, e.g. biological N<sub>2</sub> fixation (Liu et al., 2011), water drainage (Bruelle et al., 2017), N<sub>2</sub>O emissions or nitrate leaching (Plaza-Bonilla et al., 2015).

The *STICS* model is a generic dynamic soil-crop model that has been used to assess the agronomic and environmental performances of crop rotations and to support the design of innovative cropping systems in a wide range of agro-pedo-climatic conditions (Coucheney et al., 2015). For example, Bécél et al. (2015) assessed the impact of various integrated weed management strategies on nitrate leaching. Constantin et al. (2015a, 2015b) determined the optimal date of emergence and destruction of cover crops to reduce nitrate leaching. Jégo et al. (2010) analyzed the effect of fertilization, irrigation strategies and climate variability on nitrate leaching in sugar beet/potato cropping systems. So far, spring pea (*Pisum Sativum* L.) and soybean (*Glycine max* (L.) Merr.) are the only grain legume calibrated for *STICS* (Corre-Hellou et al., 2009; Jégo et al., 2010). *STICS* can simulate N sharing between companion crops and N carry over effects on following crops, but all the involved crops must be calibrated for *STICS*. Given the potential of grain legume to improve cropping system sustainability in Europe (Watson et al., 2017) and the potential of crop models to help with the design of innovative and resilient cropping systems including legumes, there is a need to broaden the number of grain legume crops calibrated and evaluated with the *STICS* cropping system model.

Faba bean (*Vicia Faba* L.) is the second most widely grown European grain legumes (Watson et al., 2017). Its proportion of crop N derived from atmospheric N<sub>2</sub> is the greatest among major grain legumes cultivated in Europe (153 kg N ha<sup>-1</sup> on average) (Jensen et al., 2010). It is a high-biomass crop leaving more residues and giving a greater nitrogen-related yield effect on the subsequent crop compared with low biomass crop like chickpea and lentil for example (Watson et al., 2017). However faba bean, like other legume crops, can suffer from strong yield and N<sub>2</sub> fixation reduction due to abiotic factors (Bishop et al., 2016; Khan et al., 2010; López-Bellido et al., 2011) and pests and diseases (Stoddard et al., 2010).

Winter faba bean experiments were conducted in southwestern

France over several growing seasons (2002–2015) with limited pest, disease and weeds pressure. This dataset provides a rich basis to adapt *STICS* to faba bean and analyse the capacity of the model to unravel the role of abiotic factors in driving yield and N<sub>2</sub> fixation variability. Our first objective was to calibrate and assess the performance of *STICS* to simulate N acquisition by soil mineral uptake and biological N<sub>2</sub> fixation, crop development and shoot biomass production, grain yield formation, N content in grain and straw, and soil water and mineral N contents along the whole profile during crop growth and after harvest. By doing so, we explored the hypothesis that the generic model (i.e. same formalism for all crops) can take into account the specificities of faba bean (N<sub>2</sub> fixation) and can be robustly calibrated for this grain legume. The second objective was to analyze the ability of *STICS* to generate a robust prediction for an independent dataset and a credible agronomic diagnosis of yield and N<sub>2</sub> fixation variability due to abiotic stresses. For this objective, our hypothesis was that the spatiotemporal adequacy of soil/crop relationships (e.g. N nutrition based on both mineral-N uptake and N<sub>2</sub> fixation) and the realism of the implemented process equations allow for an accurate prediction of abiotic stresses.

## 2. Materials and methods

### 2.1. *STICS*, a soil-crop model

#### 2.1.1. Overview of the model

The soil–crop model (as well as cropping system model) *STICS* (Brisson et al., 2009, 2002, 1998) was chosen for its agro-environmental purpose, crop genericity (allowing the simulation of crop successions with the same model) and robustness. *STICS* simulates carbon, water and nitrogen dynamics in the soil–crop system on a daily time step over the crop cycle. Unlike crop models that combine crop modules with different structures, e.g. *DSSAT* (Decision Support System for Agrotechnology Transfer, Jones et al., 2003) or *APSIM* (Agricultural Production Systems sIMulator, Keating et al., 2003), *STICS* uses a common and generic structure for different crop species. Species-specific sets of parameters define the differences between species. *STICS* has been adapted and tested for different type of crops (e.g. winter wheat, maize, barley, sunflower, rapeseed, grassland, tomato, grapevines) (Coucheney et al., 2015).

Inputs to the model include daily weather information (minimum and maximum temperature, rainfall, wind and global radiation), soil characteristics (per-layer initial soil water and mineral N content, field capacity and wilting point) and crop/soil management (date and depth of planting, planting density, soil tillage, residues management, organic and mineral fertilisation, irrigation).

Crops are defined by species parameters, ecophysiological options (e.g. effect of photoperiod and/or cold requirements on crop phenology, potential radiation use efficiency) and cultivar specific parameters (e.g. flowering precocity, maximum number of grains per m<sup>2</sup>). Crop temperature (calculated from weather variables) and photoperiod drive crop phenology. The model dynamically simulates (i) development of the root system that takes up N and water according to root density over the whole soil profile and (ii) the establishment of the canopy that transpires water and intercepts light to produce the crop biomass.

Two processes account for yield formation in the model: (i) setting of grain number per square meter and (ii) accumulation of biomass in grain. The number of grains before the start of grain filling is a function of: (i) the number of days before the beginning of grain filling during which grain number is determined (crop parameter), (ii) the mean canopy growth rate during this phase (calculated by the model), (iii) the sensitivity of grain onset to growth conditions (crop parameter) and (iv) the maximum number of grain (crop parameter). Dry matter accumulation in grains is calculated by applying a dynamic harvest index (grain to shoot biomass ratio) to the dry weight of the plant (Amir and Sinclair, 1991). This harvest index increases with time from the beginning of grain filling to physiological maturity, the rate of increase

**Table 1**  
Climate and soil characteristics (averaged across growing seasons and experimental fields) in the two experimental sites (standard error in brackets).

| Site      | Number of growing seasons | November–July                                           |                                         |                          |    |             | Number of experimental fields | Clay content (0–30 cm) (%) | pH (0–30 cm) | Maximum rooting depth (cm) | Gravimetric moisture at field capacity (%) | Gravimetric moisture at wilting point (%) | Maximum available water <sup>a</sup> (mm to maximum rooting depth) |
|-----------|---------------------------|---------------------------------------------------------|-----------------------------------------|--------------------------|----|-------------|-------------------------------|----------------------------|--------------|----------------------------|--------------------------------------------|-------------------------------------------|--------------------------------------------------------------------|
|           |                           | Cumulative global solar radiation (MJ m <sup>-2</sup> ) | Cumulative growing degree days (°C day) | Cumulative rainfall (mm) |    |             |                               |                            |              |                            |                                            |                                           |                                                                    |
| Auch      | 7                         | 3354 (64.8)                                             | 3083 (59.1)                             | 542 (47.2)               | 11 | 31.8 (1.46) | 8.4 (0.03)                    | 70 (4.9)                   | 17.4 (0.45)  | 12.0 (0.27)                | 64 (4.8)                                   |                                           |                                                                    |
| Auzeville | 7                         | 3561 (30.4)                                             | 3412 (49.1)                             | 528 (42.8)               | 12 | 25.9 (0.97) | 7.4 (0.15)                    | 135 (4.5)                  | 19.4 (0.43)  | 10.2 (0.23)                | 167 (8.5)                                  |                                           |                                                                    |

\* Average across 0–30 cm, 30–60 cm, 60–90 cm and 90–120 cm soil layers.

\*\* Soil water content at field capacity minus soil water content at wilting point.

being a crop parameter. A maximum final harvest index (crop parameter) prevent unrealistic remobilization levels. A maximum grain yield corresponding to the simulated number of grains multiplied by the maximum weight of one grain (crop parameter) is computed to avoid unrealistic yield level.

The model also accounts for soil water and N dynamics. Net N mineralization from crop residues and soil organic matter (affected by clay and CaCO<sub>3</sub> content in the topsoil and governed by soil moisture and temperature), nitrification (affected by pH in the topsoil), nitrate leaching and ammonia and nitrous oxide gaseous emissions are daily simulated as well as vertical water drainage when field capacity is exceeded.

STICS also simulates N acquisition by N<sub>2</sub> fixation for legumes. Nodule formation is a function of soil thermal time and sets the potential fixation. Actual N<sub>2</sub> fixation then depends on shoot growth rate (C limitation for N<sub>2</sub> fixation), soil temperature, soil moisture, and a nitrate inhibition effect linked with nitrate-N concentration in the nodulation layer (see Section 2.1.2).

The equations governing physical and biological processes occurring in the soil/crop system are based on a unique set of general parameters. Model parameters and equations describing yield formation are of particular interest in this study and are described in details in Table S2. An exhaustive description of inputs, equations and default parameter values of the STICS model is given in Brisson et al. (2008). Bergez et al. (2014) give an overview of the latest plant and soil process-based developments of the STICS model, and Coucheney et al. (2015) detailed the STICS version 8.5 used in this study.

#### 2.1.2. Stress factors

Physical and biological processes are successively addressed in two types of equations: 1) the calculation of the potential flux or rate of the process and 2) the effect of the environmental or crop related limiting factors. Four stress factors can indirectly affect grain yield through plant growth. The first is a water stress factor that affects radiation use efficiency and plant transpiration, and is computed as the ratio of actual evapotranspiration over potential evapotranspiration. Potential transpiration is function of Leaf Area Index (LAI) and daily potential Penman evapotranspiration, as calculated by the model. Actual transpiration is calculated as the minimum between water soil supply and maximal crop demand. The second stress factor is a Nitrogen stress that affects i) LAI increase, ii) radiation use efficiency and iii) possibly senescence. The N stress index is computed as the ratio of actual Nitrogen concentration over Critical crop N concentration. Critical crop N concentration is the minimum N concentration that allow to maximise biomass production (Lemaire and Gastal, 1997). Actual N uptake is calculated as the minimum between soil supply and maximal crop demand. The third stress is a heat reduction factor on radiation use efficiency that increases when daily average crop temperature exceeds optimal crop temperature and is maximal when daily average crop temperature exceeds maximal crop temperature. The fourth is a heat stress factor that stops grain yield when daily average crop temperature exceeds maximal temperature for grain filling.

For legumes, STICS also calculates water, heat, anoxia and nitrogen stress factors for N<sub>2</sub> fixation. The water deficit factor for N<sub>2</sub> fixation is the proportion of soil layers in the nodulation area (defined by the crop parameter “*profnod*”, set to 30 cm in our study) for which moisture is above wilting point. The heat stress factor for N<sub>2</sub> fixation increases and is maximal when soil temperature in nodulation zone exceeds optimal and maximal temperature thresholds. Nitrogen stress for N<sub>2</sub> fixation increases and is maximal when the mean amount of mineral nitrogen in the nodulation area exceeds an optimal or maximal N concentration threshold. Eventually, anoxia stress factor for N<sub>2</sub> fixation increases with the proportion of soil layers in anaerobic conditions.

Stress factors are computed daily and vary between 0 (complete stress) and 1 (no stress). Model parameters and equations used to compute water stress factor and water deficit factor for N<sub>2</sub> fixation are

of particular interest in this study and are described in details in Table S1. The exhaustive description of the equations and parameters governing the stresses definition can be found in Brisson et al. (2008).

## 2.2. Experimental data

### 2.2.1. Sites and experimental design

The experimental data was collected in south of France at the experimental sites of INRA in Auzeville (43°31'N 1°28'E), and of CREAB-MP in Auch (43°38'N 0°36'E) during seven growing seasons in each site. In both sites, the climate is oceanic temperate under mediterranean influence and characterized by summer droughts and cool wet winters. Cumulative global radiation, growing degree-days and rainfall during crop growth (November–July), averaged across growing seasons, were similar in Auch and Auzeville (Table 1). Experimental plots were in the valley on deep (i.e. average maximum rooting depth of 135 cm) clay-loamy soils in Auzeville and on the hillside on shallow (i.e. average maximum rooting depth of 70 cm) clay loamy soils in Auch. As a result, average maximum available water to maximum rooting depth (i.e. soil water content at field capacity minus soil water content at wilting point) was higher in Auzeville (167 mm) than in Auch (64 mm) (Table 1).

Cropping systems strongly differed between experimental sites. In Auzeville, the data originated from two distinct types of experiments. The first type was annual experiments with replicated plots (1.62 m wide × 13 m long) of sole faba bean (see Bedoussac and Justes (2010) and Kammoun (2014) for a detailed description of experimental design). The second type was a long-term cropping system experiment established in 2003 on 200 × 30 m plots with different three-years rotations varying in the frequency of grain legumes and the use or not of cover crops (Plaza-Bonilla et al., 2017). Appropriate herbicide and pesticides were applied to control weeds, pests and diseases and no N fertilizer was applied on legumes. In Auch, the data originated from an organic long-term soil fertility monitoring experiment (2002–2013) with 50 × 50 m plots with Faba bean – Oat or wheat – Sunflower rotations (Colomb et al., 2013). This site was managed without mineral fertiliser, herbicides and pesticides. As a result, only the plots where the monitoring revealed negligible pests, disease and weed pressure were considered. In both sites, the land was plowed prior to sowing and only in 2007 in Auzeville small amounts of irrigation water (20–50 mm) was applied after sowing (Table 2) to ensure satisfactory crop establishment.

The considered faba bean cultivars were Castel and Irena (Table 2). These winter-type cultivars sown between November and January have similar phenology and yield potential (Flores et al., 2012). Site, year, and management factors (cultivar, crop density, incorporation of a cover crop before planting and sowing date) defined 35 Site-Year-Management units (Table 2). Site-year-management units covering a broad range of soil types, growing seasons and management situations (He et al., 2017) were chosen for the calibration dataset (Table 2). The remaining units, differing in growing season and/or management (compared with calibration dataset) formed the evaluation dataset (Table 2).

### 2.2.2. Measurements

Faba bean development stages were recorded in the annual experiments in Auzeville following BBCH scale (Meier, 2001), namely (i) BBCH 09: 50% of shoot have emerged through soil surface (ii) BBCH 70: first pods have reached final length (corresponding to “drp” stage in STICS, i.e. the beginning of grain filling) and (iii) BBCH 89: Fully ripe: nearly all pods dark, seeds dry and hard. Additionally, interpolation of discontinuous LAI observations allowed for determining the date of the end of juvenile phase (*amf* stage in STICS, corresponding to maximum growth rate of LAI). The interpolation was carried out with the tool integrated in STICS that uses a functional equation to differentiate LAI growth (logistic equation) and senescence (exponential equation), the

difference between the two functions giving the green LAI (see Casa et al. (2012) for detailed equations and procedure).

In each experimental field, soil texture and pH were measured over the whole soil profile (0–120 cm). Soil mineral N and water content were measured by layers of 30 cm three times per year: (1) end of autumn prior to sowing, (2) mid-winter and (3) at harvest. Crop variables such as LAI, above-ground biomass, fixed N<sub>2</sub> and total accumulated N, grain yield and grain N content were measured from 1 to 5 times depending on site, year and experiment (Table 2): (1) in mid-March i.e. wheat ‘1 cm ear’ stage, (2) in mid-April, i.e. legume flowering, (3) in mid-May i.e. wheat flowering, (4) in early June i.e. wheat ripening, and (5) in early July, i.e. legume maturity. At each measurement, six central rows 0.5 m long (annual experiments in Auzeville), six m<sup>2</sup> (long term cropping system experiment in Auzeville) and nine subplots of two rows 1 m long (in Auch) were harvested by cutting plants at the soil surface. In the annual experiments in Auzeville, a LI-3100 planimeter (LI-COR, Lincoln, NE) was used to determine green leaf area. In all experiments, plant biomass samples were dried at 80 °C for 48 h and weighted. At maturity, plants were threshed to determine grain yield and harvest index. Above-ground plant and grain N concentration were determined on finely ground plant material with the Dumas combustion method using a Leco-2000 analyzer (LECO, St. Joseph, MI, US).

In Auzeville, N<sub>2</sub> fixation ratio, i.e. the proportion of plant N derived from atmosphere (%Nd<sub>fa</sub>) was determined using the <sup>15</sup>N natural abundance method (Unkovich, 2008) with unfertilized wheat and non-legume weeds as reference plants in the annual and long-term experiments respectively. %Nd<sub>fa</sub> was calculated as follows:

$$\%Nd_{fa} = 100 \times \frac{\delta^{15}N_{ref} - \delta^{15}N_{fababeau}}{\delta^{15}N_{ref} - \beta} \quad (1)$$

where  $\delta^{15}N$  is the deviation from the international standard of atmospheric N<sub>2</sub> (0.3663‰ <sup>15</sup>N) for faba bean ( $\delta^{15}N_{fababeau}$ ) and for the reference crop ( $\delta^{15}N_{ref}$ ), and  $\beta$  is the  $\delta^{15}N$  of shoots of faba bean fully dependent upon N<sub>2</sub> fixation, assumed equal to −0.63‰ (Unkovich, 2008). The amount of N<sub>2</sub> fixed was calculated as the product of %Nd<sub>fa</sub> and total above-ground plant N. Moreover, dynamics of maximum rooting depth of faba bean was monitored in 2013 during the annual experiment in Auzeville (Table 2) using a rotary scanner-based minirhizotron system (CI-600, CID Bio-science, Camas, WA, USA) six times during crop growth (16 and 28 March, 12 April, 6 May, 3 June, 17 July).

## 2.3. Calibration procedure

The calibration consisted in choosing the formalisms and corresponding parameters involved in the simulation of crop phenology, leaf area and root system expansion, biomass accumulation, biological nitrogen fixation, nitrogen uptake and grain yield elaboration.

The parameterization procedure included three steps (Fig. 1). The first step was a literature review to select the options offered by the model to simulate the different crop processes and determine existing parameter values. The second step corresponded to the determination of parameters from experimental data by direct measurement or observation (phenology). The last step consisted in a mathematical optimisation based on the approach developed by Guillaume et al. (2011), including nine stages representing the different key processes in the simulation of the final outputs of the model (Table 3). For each stage, sensitivity analysis (Ruget et al., 2002) and expert knowledge were used to select the parameters having the strongest impact on model outputs. In total, 29 crop parameters were mathematically optimised (Table 3). Stage-specific target variables were selected: (i) dates of key development stages (emergence, end of juvenile phase, beginning of grain filling and physiological maturity), (ii) dynamic variables (Leaf Area Index, above-ground biomass, fixed N and total plant N, overall soil moisture content and overall soil N content) and (iii) variables at

**Table 2**

Description of the Site-Year-Management units in the calibration and evaluation datasets. LAI: Leaf Area Index, AGPN: above-ground plant N, AGB: above-ground biomass, GY: grain yield, GNC: grain N content, SN: Soil N content over the entire profile, SWC: soil water content over the entire profile.

| Dataset     | Site      | Year (harvest) | Management |                                            |                            |             |                                      | Number of measurements during growing season |         |      |     |    |     |            |    |     |
|-------------|-----------|----------------|------------|--------------------------------------------|----------------------------|-------------|--------------------------------------|----------------------------------------------|---------|------|-----|----|-----|------------|----|-----|
|             |           |                | Cultivar   | Density (Number of plant m <sup>-2</sup> ) | Cover crop before planting | Sowing date | Dates and amount of irrigation water | LAI                                          | Fixed N | AGPN | AGB | GY | GNC | Root depth | SN | SWC |
| Calibration | Auzeville | 2007           | Castel     | 30                                         | No                         | 7 November  | 7 November - 20mm                    | 4                                            | 1       | 5    | 5   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 15                                         | No                         | 7 November  | 7 November - 20mm                    | 4                                            | 1       | 5    | 5   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2010           | Castel     | 30                                         | No                         | 19 November | 0                                    | 4                                            | 2       | 5    | 5   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 15                                         | No                         | 19 November | 0                                    | 4                                            | 2       | 5    | 5   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2011           | Irena      | 30                                         | No                         | 2 December  | 14 April; 16 May - 34;49 mm          | 4                                            | 4       | 5    | 5   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Irena      | 30                                         | No                         | 2 December  | 0                                    | 0                                            | 2       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2012           | Castel     | 30                                         | No                         | 22 November | 0                                    | 0                                            | 1       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 15                                         | No                         | 22 November | 0                                    | 0                                            | 1       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2013           | Irena      | 30                                         | No                         | 20 November | 0                                    | 4                                            | 1       | 1    | 4   | 1  | 1   | 6          | 2  | 2   |
|             |           |                | Irena      | 30                                         | No                         | 23 November | 0                                    | 0                                            | 0       | 0    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2014           | Irena      | 15                                         | No                         | 23 November | 0                                    | 4                                            | 1       | 1    | 4   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Irena      | 30                                         | No                         | 23 November | 0                                    | 0                                            | 0       | 0    | 3   | 1  | 1   | 0          | 2  | 2   |
|             | Auch      | 2002           | Castel     | 30                                         | No                         | 17 December | 0                                    | 0                                            | 0       | 1    | 1   | 1  | 1   | 0          | 1  | 1   |
|             |           |                | Castel     | 30                                         | No                         | 17 December | 0                                    | 0                                            | 0       | 1    | 1   | 1  | 1   | 0          | 1  | 1   |
|             |           | 2003           | Castel     | 30                                         | No                         | 20 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 30                                         | No                         | 20 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2004           | Castel     | 30                                         | No                         | 25 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 30                                         | No                         | 12 January  | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2005           | Castel     | 30                                         | No                         | 12 January  | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 30                                         | No                         | 23 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2007           | Castel     | 30                                         | No                         | 1 December  | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 30                                         | No                         | 1 December  | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2010           | Castel     | 30                                         | No                         | 1 December  | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 30                                         | No                         | 11 December | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2010           | Castel     | 30                                         | No                         | 11 December | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Castel     | 30                                         | No                         | 11 December | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | Total          |            |                                            |                            |             |                                      | 28                                           | 15      | 55   | 64  | 22 | 22  | 6          | 42 | 42  |
| Evaluation  | Auzeville | 2012           | Castel     | 30                                         | Yes                        | 22 November | 0                                    | 0                                            | 1       | 3    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2013           | Irena      | 30                                         | Yes                        | 23 November | 0                                    | 0                                            | 0       | 0    | 3   | 1  | 0   | 0          | 2  | 2   |
|             |           | 2014           | Mix*       | 30                                         | Yes                        | 13 December | 0                                    | 0                                            | 3       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2015           | Mix*       | 30                                         | No                         | 13 December | 0                                    | 0                                            | 2       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Mix*       | 30                                         | Yes                        | 13 January  | 0                                    | 0                                            | 1       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2015           | Mix*       | 30                                         | No                         | 13 January  | 0                                    | 0                                            | 1       | 3    | 3   | 1  | 1   | 0          | 2  | 2   |
|             | Auch      | 2003           | Irena      | 30                                         | No                         | 20 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Irena      | 30                                         | No                         | 21 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2013           | Irena      | 30                                         | No                         | 22 November | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Irena      | 30                                         | No                         | 17 December | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | 2013           | Irena      | 30                                         | No                         | 17 December | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           |                | Irena      | 30                                         | No                         | 17 December | 0                                    | 0                                            | 0       | 2    | 2   | 1  | 1   | 0          | 2  | 2   |
|             |           | Total          |            |                                            |                            |             |                                      | 0                                            | 8       | 29   | 31  | 13 | 12  | 0          | 26 | 26  |

\* Irena, Organdi and Diver.




Fig. 1. Summary of the generic method used for calibration and evaluation of faba bean with the STICS soil/crop model and data used for this study.

**Table 3**  
Description of the nine stages undertaken for parameter optimisation: parameters optimised, minimum and maximum values and target variable. Dd = degree (°C) day.

| Stage | Simulated process | Parameter   |                                                                          |                                                     | Target variable |         |                            | Unit                                    | Description                             | Unit                   |
|-------|-------------------|-------------|--------------------------------------------------------------------------|-----------------------------------------------------|-----------------|---------|----------------------------|-----------------------------------------|-----------------------------------------|------------------------|
|       |                   | Acronym     | Description                                                              | Unit                                                | Minimum         | Maximum | Acronym                    |                                         |                                         |                        |
| 1     | Emergence         | belong      | parameter of coleoptile elongation curve                                 | Dd <sup>-1</sup>                                    | 0.005           | 0.04    | ilev                       | date of emergence                       | date of emergence                       | julian day             |
| 2a    | Crop development  | celong      | parameter of plantlet elongation curve                                   | –                                                   | 1               | 10      | ilev                       | date of emergence                       | date of emergence                       | julian day             |
|       |                   | sensiphot   | index of photoperiod sensitivity (1 = insensitive)                       | –                                                   | 0               | 1       | iamf                       | date of end of juvenile phase           | date of end of juvenile phase           | julian day             |
|       |                   | jvc         | number of vernalising days                                               | days                                                | 0               | 70      | iamf                       | date of end of juvenile phase           | date of end of juvenile phase           | julian day             |
|       |                   | phobase     | basal threshold controlling photoperiod slowing effect                   | hours                                               | 5               | 12      | iamf                       | date of end of juvenile phase           | date of end of juvenile phase           | julian day             |
|       |                   | phosat      | saturating threshold controlling photoperiod slowing effect              | hours                                               | 6               | 14      | iamf                       | date of end of juvenile phase           | date of end of juvenile phase           | julian day             |
| 2b    | Crop development  | stlevamf    | cumulative thermal time between emergence and end of juvenile phase      | Dd                                                  | 100             | 1000    | iamf                       | date of end of juvenile phase           | date of end of juvenile phase           | julian day             |
|       |                   | stamflax    | cumulative thermal time between end of juvenile phase and maximum LAI    | Dd                                                  | 100             | 1000    | ilax                       | date when LAI is maximum                | date when LAI is maximum                | julian day             |
|       |                   | stlevdrp    | cumulative thermal time between emergence and beginning of grain filling | Dd                                                  | 100             | 1500    | idrp                       | date of beginning of grain filling      | date of beginning of grain filling      | julian day             |
| 3a    | Leaves            | dlaimaxbrut | maximum rate of the setting up of LAI                                    | m <sup>2</sup> plant <sup>-1</sup> Dd <sup>-1</sup> | 0.0005          | 0.05    | lai(n)                     | Leaf Area index                         | Leaf Area index                         | –                      |
|       |                   | adens       | interplant competition parameter                                         | –                                                   | –2              | 0       | lai(n)                     | Leaf Area index                         | Leaf Area index                         | –                      |
|       |                   | bdens       | minimal density above which interplant competition starts                | plant m <sup>-2</sup>                               | 1               | 30      | lai(n)                     | Leaf Area index                         | Leaf Area index                         | –                      |
| 4     | Root growth       | durvief     | minimal lifespan of an adult leaf                                        | –                                                   | 10              | 500     | lai(n)                     | Leaf Area index                         | Leaf Area index                         | –                      |
|       |                   | croirac     | elongation rate of the root apex                                         | cm Dd <sup>-1</sup>                                 | 0.005           | 0.5     | zrac                       | Rooting depth                           | Rooting depth                           | cm                     |
|       |                   | draclong    | maximum rate of root length production per plant                         | cm plant <sup>-1</sup> Dd <sup>-1</sup>             | 1               | 1000    | HR(1), HR(2), HR(3), HR(4) | Soil water content in layer 1,2,3,4 (%) | Soil water content in layer 1,2,3,4 (%) | –                      |
| 5a    | Shoot growth      | efcroijuv   | maximum radiation use efficiency during the juvenile phase               | g MJ <sup>-1</sup>                                  | 1               | 4       | masec                      | Above ground biomass                    | Above ground biomass                    | t ha <sup>-1</sup>     |
|       |                   | efcroirepro | maximum radiation use efficiency during the grain filling phase          | g MJ <sup>-1</sup>                                  | 2               | 5       | masec                      | Above ground biomass                    | Above ground biomass                    | t ha <sup>-1</sup>     |
|       |                   | efcroiveg   | maximum radiation use efficiency during the vegetative stage             | g MJ <sup>-1</sup>                                  | 2               | 5       | masec                      | Above ground biomass                    | Above ground biomass                    | t ha <sup>-1</sup>     |
| 5b    | Shoot growth      | abscission  | fraction of senescent leaves falling to the soil                         | –                                                   | 0               | 100     | masec                      | Above ground biomass                    | Above ground biomass                    | t ha <sup>-1</sup>     |
|       |                   | ratosen     | fraction of senescent biomass (relative to total biomass)                | –                                                   | 0               | 1       | masec                      | Above ground biomass                    | Above ground biomass                    | t ha <sup>-1</sup>     |
|       |                   | fixmaxgr    | maximal N symbiotic fixation rate per unit of grain growth rate          | kg t <sup>-1</sup>                                  | 0               | 50      | Qfix                       | Above ground fixed plant N              | Above ground fixed plant N              | kg ha <sup>-1</sup>    |
| 6     | Nitrogen fixation | fixmaxveg   | maximal N symbiotic fixation rate per unit of vegetative growth rate     | kg t <sup>-1</sup>                                  | 0               | 50      | Qfix                       | Above ground fixed plant N              | Above ground fixed plant N              | kg ha <sup>-1</sup>    |
|       |                   | Knabs2      | affinity constant of N uptake by roots for the low uptake system         | μmole L <sup>-1</sup>                               | 4000            | 40000   | QNplant                    | Total Above ground plant N              | Total Above ground plant N              | kg ha <sup>-1</sup>    |
|       |                   | Vmax2       | maximum specific N uptake rate with the high affinity transport system   | μmole cm <sup>-1</sup> h <sup>-1</sup>              | 0.002           | 0.1     | QNplant                    | Total Above ground plant N              | Total Above ground plant N              | kg ha <sup>-1</sup>    |
| 8a    | Yield formation   | cgrain      | slope of the relationship between grain number and growth rate           | grains g <sup>-1</sup> day                          | 0.01            | 1       | chargefruit                | Number of filling grains                | Number of filling grains                | grains m <sup>-2</sup> |
|       |                   | nbjgrain    | number of days used to compute the number of viable grains               | days                                                | 5               | 40      | chargefruit                | Number of filling grains                | Number of filling grains                | grains m <sup>-2</sup> |
|       |                   | cgrainv0    | number of grains produced when growth rate is zero                       | grains m <sup>-2</sup>                              | 0               | 15000   | chargefruit                | Number of filling grains                | Number of filling grains                | grains m <sup>-2</sup> |
| 8b    | Yield formation   | vitircarb   | rate of increase of the C harvest index versus time                      | g grain g <sup>-1</sup> day <sup>-1</sup>           | 0.001           | 0.02    | mafruit                    | Grain yield                             | Grain yield                             | t ha <sup>-1</sup>     |
|       |                   | tmaxremp    | maximal temperature above which grain filling stops                      | °C                                                  | 10              | 40      | mafruit                    | Grain yield                             | Grain yield                             | t ha <sup>-1</sup>     |
|       |                   | vitirazo    | rate of increase of the N harvest index versus time                      | g grain g <sup>-1</sup> d <sup>-1</sup>             | 0.001           | 0.06    | QNgrain                    | Grain N content                         | Grain N content                         | kg ha <sup>-1</sup>    |

crop maturity (yield components, grain yield and grain N content) (Table 3). The calibration stages were carried out by using the software package *OptimiSTICS* (Wallach et al., 2011) and the methodology detailed by Buis et al. (2011). Model error was computed as the difference between measured and simulated values for a given target variable at a given date. The goodness of fit  $C_v$  was obtained following the formula:

$$C_v = \sum_s \frac{1}{n_s} \sum_i [O_{s,i} - P_{s,i}(\theta)]^2 \quad (2)$$

Where  $v$  is the target variable,  $s$  is the site-year-management unit and  $n_s$  is the number of measurements in  $s$ ,  $O_{s,i}$  is the measured value in site-year-management  $s$  on the  $i$ th date and  $P_{s,i}(\theta)$  is the corresponding simulated value with the parameter set  $\theta$ .

This average squared error for a given target variable was minimised using a simplex algorithm. For each optimised parameter, lower and upper limits corresponding to reasonable physiological values were fixed based on literature and expert knowledge (Table 3). The optimised parameter values at a given stage were the default values of the next steps. The initial plant parameters before optimisation were taken from the spring pea plant file (Corre-Hellou et al., 2009). Given the similarities in crop functioning and yield potential observed in our experiments between the two studied faba bean cultivars (Irena and Castel), we did not calibrate specific cultivar parameters.

Soil analysis informed the input parameters required for soil in *STICS*. Moisture at field capacity and wilting point were first obtained using pedo-transfer functions (Saxton and Rawls, 2006) and also based on laboratory measurements on sieved soil for field capacity, and then adjusted by trial and error to minimize the error between simulated and observed soil water content in mid-winter. The recorded crop managements (Table 2) were used to create management files used for each simulation. Initial soil mineral nitrogen (nitrate ammonium) and water content were measured for each experiment and set between 13 and 120 kg N ha<sup>-1</sup> and 162 and 342 mm respectively, depending on the different Site-year-management units (Table 2).

#### 2.4. Model evaluation

Model performance (with optimised parameter set) was evaluated graphically and quantified by calculating Mean Bias Error (MBE) and its relative value (rMBE), Model Efficiency (EF), Root Mean Square Error (RMSE) and its relative value (rRMSE):

$$MBE = \frac{1}{n} \sum_{i=1}^n (P_i - O_i) \quad (3)$$

$$rMBE = \frac{MBE}{\bar{O}} \times 100 \quad (4)$$

$$EF = 1 - \frac{\sum_{i=1}^n (O_i - P_i)^2}{\sum_{i=1}^n (O_i - \bar{O})^2} \quad (5)$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (O_i - P_i)^2} \quad (6)$$

$$rRMSE = \frac{RMSE}{\bar{O}} \times 100 \quad (7)$$

where  $O_i$  and  $P_i$  are the observed and simulated values for the  $i$ th measurement,  $n$  is the number of observations and  $\bar{O}$  is the mean of the observed values.

MBE and rMBE indicate whether the model under/overestimate a given variable, and gives the absolute and relative magnitude and direction of the bias. EF is based on the comparison of the simulation model with a constant model (i.e. the average of measured data) and ranges between  $-\infty$  and one. Negative EF values indicate that the constant model performs better than or as well as the simulation model. A value of one indicate a perfect model. EF is a good indicator to compare

model simulations with two different sets of parameters (e.g. before/after calibration) against the same dataset. It is however of little value to compare two simulations with the same set of parameters against two different datasets, because it depends on the variance of the observed dataset. RMSE and rRMSE are the model absolute and relative prediction error. The combined analysis of these four indicators give a robust assessment of model accuracy. Similarly to Beaudoin et al. (2008) and Constantin et al. (2015a, 2015b), model predictions were judged satisfactory when EF value were greater than 0.5 and rMBE lower than 10%. A summary of the calibration and evaluation procedure can be found in Fig. 1.

#### 2.5. Stress factors and variability in grain yield and total N<sub>2</sub> fixed

Stress factors calculated by *STICS* (see Section 2.1.2 and Table S1) were averaged over periods ranging from -60 to + 60 days before/after the beginning of grain filling. For each period, the Pearson correlation coefficient ( $R^2$ ) between observed grain yield (total fixed N<sub>2</sub> respectively) and stress factor affecting grain yield (N<sub>2</sub> fixation respectively) over the period was computed. The period corresponding to the maximum  $R^2$  was determined. For this analysis, calibration and evaluation datasets were pooled.

### 3. Results

#### 3.1. Parameter values obtained from literature, experimental data and optimisation

Eight parameters values were obtained from literature (Table 4). Faba bean shows a long day response to photoperiod (Ellis et al., 1988) and needs vernalisation (Patrick and Stoddard, 2010) so we activated these physiological options in the faba bean plant file. Faba bean base temperature for crop development ( $tdmin$ ) was set to 0 °C (Tribouillois et al., 2016). Because there was no available data in the literature, base and optimal temperature for photosynthesis ( $temin$  and  $teopt$ ) were approximated from those for emergence rate (Tribouillois et al., 2016) (Table 4), based on the assumption that cardinal temperatures for developmental processes are close for a given species (e.g. Parent and Tardieu, 2012). Cardinal temperatures for nodulation ( $tempnod1$ ,  $tempnod2$ ,  $tempnod3$ ,  $tempnod4$ ) were obtained from Boote et al. (2002) (Table 4).

Four parameters values were obtained using direct in-field measurement (Table 4 and Fig. 2). Maximum and minimum number of grain per square meter ( $nbgrmax$  and  $nbgrmin$ ), maximum grain weight ( $pgrainmaxi$ ) and maximum harvest index ( $irmax$ ) were set to the maximum or minimum measured values in the experiments in Auzeville and Auch with an added or subtracted 10% respectively to reach assumed potential values.

The other 30 parameters values were set according to the result of the sequential optimisation with *OptimiSTICS* using measured variables (Table 4).

#### 3.2. Agreement between simulated and observed values

##### 3.2.1. Crop development and LAI

Calibration led to satisfactory prediction of the dates of emergence, end of juvenile phase, beginning of grain filling and physiological maturity. Relative bias was low (0–10%) with fair EF values (Fig. 3), indicating the relevance of calculating crop development based on vernalisation and photoperiodic and thermal time-course formalisms. Simulated LAI values for the calibration dataset were close to observed values with a low relative bias of 1% and a satisfactory EF of 0.69. Prediction error was however great due to a bias of a few days in the senescence dynamic that had large consequences on the daily agreement between observed and simulated LAI values (rRMSE = 56%) (Fig. 4a).

**Table 4**Values of faba bean parameter as calibrated in the *STICS* crop model for experiments in Auch and Auzéville, France.

| Parameter         |                |                                                                         |            |                            |
|-------------------|----------------|-------------------------------------------------------------------------|------------|----------------------------|
| Process           | Acronym        | Description                                                             | Value      | Source                     |
| Emergence         | tdmin          | basal temperature for crop development                                  | 0          | Tribouillois et al. (2016) |
|                   | belong         | parameter of coleoptile elongation curve                                | 0.0200     | mathematical optimisation  |
|                   | celong         | parameter of plantlet elongation curve                                  | 6.20       | mathematical optimisation  |
| Crop development  | sensiphot      | index of photoperiod sensitivity (1 = insensitive)                      | 0.6        | mathematical optimisation  |
|                   | jvc            | number of vernalising days                                              | 3          | mathematical optimisation  |
|                   | phobase        | basal photoperiod controlling photoperiod slowing effect                | 5.7        | mathematical optimisation  |
|                   | phosat         | saturating photoperiod controlling photoperiod slowing effect           | 12         | mathematical optimisation  |
|                   | stlevamf       | cumulative thermal time between emergence and end of juvenile phase     | 518        | mathematical optimisation  |
|                   | stamflax       | cumulative thermal time between end of juvenile phase and maximum LAI   | 700        | mathematical optimisation  |
| Leaves            | stlevdrp       | cumulative thermal time between emergence and begining of grain filling | 1180       | mathematical optimisation  |
|                   | dlaimaxbrut    | maximum rate of the setting up of LAI                                   | 0.0025     | mathematical optimisation  |
|                   | adens          | interplant competition parameter                                        | −0.50      | mathematical optimisation  |
|                   | bdens          | minimal density above which interplant competition starts               | 1          | mathematical optimisation  |
|                   | durvieF        | maximal lifespan of an adult leaf                                       | 220        | mathematical optimisation  |
|                   | croirac        | elongation rate of the root apex                                        | 0.07       | mathematical optimisation  |
| Root growth       | draclong       | maximum rate of root length production per plant                        | 810        | mathematical optimisation  |
| Shoot growth      | efcroijuv      | maximum radiation use efficiency during the juvenile phase              | 1          | mathematical optimisation  |
|                   | efcroirepro    | maximum radiation use efficiency during the grain filling phase         | 3.5        | mathematical optimisation  |
|                   | efcroireveg    | maximum radiation use efficiency during the vegetative stage            | 2.3        | mathematical optimisation  |
|                   | abscission     | fraction of senescent leaves falling to the soil                        | 1          | mathematical optimisation  |
|                   | ratiosen       | fraction of senescent biomass (relative to total biomass)               | 0.75       | mathematical optimisation  |
|                   | temin          | basal temperature for photosynthesis                                    | 0          | Tribouillois et al. (2016) |
| Nitrogen fixation | teopt          | optimal temperature for photosynthesis                                  | 24         | Tribouillois et al. (2016) |
|                   | temax          | maximal temperature for photosynthesis                                  | 34         | Tribouillois et al. (2016) |
|                   | fixmaxgr       | maximal N symbiotic fixation rate per unit of grain growth rate         | 17         | mathematical optimisation  |
|                   | fixmaxveg      | maximal N symbiotic fixation rate per unit of vegetative growth rate    | 27         | mathematical optimisation  |
|                   | tempnod1,2,3,4 | cardinal temperatures driving N <sub>2</sub> fixation                   | 0,16,25,40 | Boote et al. (2002)        |
|                   | Kmabs2         | affinity constant of N uptake by roots for the low uptake system        | 33300      | mathematical optimisation  |
| Nitrogen uptake   | Vmax2          | maximum specific N uptake rate with the high affinity transport system  | 0.16       | mathematical optimisation  |
|                   | cgrain         | slope of the relationship between grain number and growth rate          | 0.0475     | mathematical optimisation  |
| Yield formation   | nbjgrain       | number of days used to compute the number of viable grains              | 9          | mathematical optimisation  |
|                   | cgrainv0       | number of grains produced when growth rate is zero                      | 0.172      | mathematical optimisation  |
| Yield formation   | vitircarb      | rate of increase of the C harvest index vs time                         | 0.04       | mathematical optimisation  |
|                   | tmaxremp       | maximal temperature above which grain filling stops                     | 33         | mathematical optimisation  |
|                   | nbgrmin        | minimum number of grains                                                | 360        | Measurement                |
|                   | nbgrmax        | maximum number of grains                                                | 1120       | Measurement                |
|                   | pgrainmaxi     | maximum weight of one grain                                             | 0.57       | Measurement                |
|                   | irmax          | maximum harvest index                                                   | 0.6        | Measurement                |
| Grain N content   | vitirazo       | rate of increase of the N harvest index versus time                     | 0.056      | mathematical optimisation  |

### 3.2.2. Above ground biomass, N<sub>2</sub> fixation and N acquisition

Predicted biomass agreed with observation throughout the growth seasons in both calibration and validation datasets ( $EF > 0.77$  and  $-7\% < \text{bias} < 0\%$ ). Prediction error was high due to senescence defoliation between grain physiological maturity and harvest that was difficult to precisely simulate (rRMSE of 41 and 31% for calibration and

evaluation respectively). There was a large variation in faba bean biomass at harvest ( $1.5\text{--}11.6 \text{ t ha}^{-1}$  and  $2.8\text{--}7.2 \text{ t ha}^{-1}$  for calibration and evaluation datasets respectively) depending on site, year and management (Fig. 4c and d) and the model was able to reproduce this behaviour.

The calibration of parameters related to N<sub>2</sub> fixation (Tables 3 and 4)




**Fig. 2.** Boxplots of number of grain per square meter (a), weigh of one grain (b) and harvest index (c) in the 35 site-year-management units of the study. The horizontal line in the box indicates the median. The height of the box represents the interquartile range. The whiskers extend to the maximum values. The horizontal dotted line correspond to the maximum measured values with an added 10% to define potential maximum values in the *STICS* model.




Fig. 3. Comparison of observed and simulated (STICS model) faba bean phenology.

allowed to reach a good agreement of simulated *versus* observed values throughout crop growth (Fig. 4e and f) ( $EF = 0.77$  with negligible bias).  $EF$  remained high and bias remained low in the evaluation indicating that the model was able to reproduce the wide range of final  $N_2$  fixed at harvest ( $62\text{--}166\text{ kg N ha}^{-1}$ , Fig. 4f).  $N$  acquired ( $N$  content in crop aerial parts) was adequately simulated with low bias both in calibration and evaluation ( $-7$  and  $2\%$  respectively) and  $EF$  values above  $0.5$  (Fig. 4g and h).

### 3.2.3. Grain yield and grain $N$ content

Observed grain yield greatly varied depending on site-year-management ( $0.42\text{--}4.7\text{ t ha}^{-1}$ , Fig. 5a). The model performed well in simulating grain yield ( $EF = 0.66$  with low relative bias) for calibration dataset. Model performance remained satisfactory during evaluation, with an  $EF$  value of  $0.65$  and a relative bias of  $-1\%$  (Fig. 5b).

Simulated grain  $N$  content was slightly underestimated (relative bias of  $-10\%$ ) in the calibration dataset with a satisfactory  $EF$  value of  $0.60$  (Fig. 5c). In the evaluation, grain  $N$  content was predicted with a low relative bias ( $3\%$ ). The  $EF$  was low ( $0.27$ ) but the range of observed values for evaluation ( $65\text{--}162\text{ kg N ha}^{-1}$ ) was narrower than for calibration ( $21\text{--}200\text{ kg N ha}^{-1}$ ).

### 3.2.4. Soil water and soil mineral $N$ contents

Soil water content over the entire profile was correctly simulated by the model, both during calibration and evaluation ( $EF > 0.70$  with small relative bias) (Fig. 5e and f). Soil  $N$  content was less adequately predicted, with small ( $0.22$ ) and negative ( $-1.42$ )  $EF$  values, in calibration and evaluation respectively (Fig. 5g and h). Bias was low in the evaluation ( $-10\%$ ), indicating that though the model was not able to reproduce precisely the differences between site, year and management, it predicted soil mineral  $N$  content within the range of the rather low observed values.

## 3.3. Impact of temporal dynamics of stress factors

With regard to grain yield, the water stress factor (*i.e.* water supply/demand ratio, see section 2.1.2 and Table S1) had the strongest explanatory power. Indeed, when averaged over a period of 6 days before the beginning of grain filling (*i.e.* the end of grain onset), it explained

78% of the observed grain yield variability in the overall dataset (Fig. 6a). The model fairly reproduced the temporal dynamic of above ground biomass (Fig. 7d–f). Water stress during the end of grain onset (Fig. 7a–c) affected the growth rate of above ground biomass (Fig. 7d–f), which in turn influenced the final number of grains (Fig. 7g–i). In the end, the model was able to simulate contrasting water stresses that occurred in different soil types (valley/hillside) for a range of climatic conditions and their final impact on the number of grain and grain yield (Fig. 7). Heat stress effect on radiation use efficiency, conversely, had a limited explanatory power in our conditions. Indeed, when averaged over a period of 26 days after the beginning of grain filling (the period with the best correlation), heat stress explained only 25% of the observed grain yield variability. The model simulated no heat stress on grain filling in the majority of the site-year-management units. There were only five days in 2002 in Auch during which crop temperature exceeded the maximal temperature above which grain filling stops.

With regard to  $N_2$  fixation, the water deficit factor had also the strongest explanatory power. Thus, when averaged over a period of 20 days after the beginning of grain filling, it explained 82% of the observed variability in total fixed  $N_2$  at harvest (Fig. 6b). Despite a limited number of within-season measurements in some situations (Fig. 8d), the model reproduced reasonably well the temporal dynamic of  $N_2$  fixation (Fig. 8c and d). In the end, the model was able to reproduce the wide range of total fixed  $N_2$  at harvest ( $62\text{--}172\text{ kg ha}^{-1}$ , see Fig. 6). The model diagnosed no stress due to anoxia in any of the cropping situations of this study. Though the model diagnosed limited heat and nitrogen stresses in some situations (value of  $0.67$  and  $0.58$  respectively) the correlation with the amount of  $N_2$  fixed was not significant for any of the periods considered.

## 4. Discussion

### 4.1. A method to adapt a generic crop model to a new crop

While the process of calibrating a new cultivar is well described, *e.g.* Jégo et al. (2010), less emphasis has been put on the description of the adaptation of a generic soil-crop model to a new crop. Previous studies mainly described a partial calibration based on a limited set of target variables, *e.g.* LAI or AGB and Soil water (Confalonieri and Bechini, 2004; Ko et al., 2009), and did not address the broad range of processes available in the model (*e.g.* root system expansion, Nitrogen uptake, yield formation). Our study shows that the generic STICS soil-crop model can be successfully adapted to a new crop, and in particular a grain legume having the specificity of  $N_2$  fixation. The adaptation was based on a parameter estimation based on measurement and literature and a sound and careful sequential optimisation of crop parameters that were not measured nor estimated using existing literature. Therefore, our work contributes to establish a generic method to calibrate a new annual crop for soil-crop models that (i) have a common structure for different crop species, (ii) are functional deterministic models considering water, C and N cycles, on a daily time step (Di Paola et al., 2016) and (iii) have an available parameter optimisation tool.

Estimating a parameter from direct measurements and literature has the advantage of not being based on the assumption that model formalism and all other parameters are reliable, as it is the case with mathematical optimisation (Grant, 2001). It should be therefore prioritised and in our approach we had 9 parameters estimated in this way (Table 4). However, this procedure is not always possible for many reasons. Firstly, field measurements of parameters is expensive and time consuming (Anothai et al., 2008). Secondly, the dataset used for model calibration may not be originally intended to cover model calibration issues (as it was the case in our study), so that most of crop model parameters are not directly measured (*e.g.* radiation use efficiency). Thirdly, the need for similar environmental conditions, varieties and exact correspondence of the measured variable to the process




**Fig. 4.** Comparison of observed and *STICS* simulated LAI, Above ground biomass (AGB),  $N_2$  fixed, and Above ground plant N (AGPN) for calibration and evaluation dataset. rMBE = relative mean bias error, EF = model efficiency, rRMSE = relative Root Mean Square Error. The black line is the 1:1 line. The dotted line represent the regression of simulated against observed values. White symbols correspond to the end of growing season.

represented by the model limit the number of parameters that can be derived from a literature review. For example, the radiation use efficiency parameters in *STICS* (*efcroijuv*, *efcroiveg* and *efcroirepro*, see Table 3) correspond to potential conditions with no stress, and the presence or absence of stress is not necessarily mentioned in the literature. Finally, there is a risk of error accumulation when extensively

relying on approximate or conflicting literature values (Wallach et al., 2011). For all these reasons, recourse to optimisation is advisable. The strength of our approach lies in the rational stepwise optimisation procedure chosen. Rather than an empirical trial and error procedure (Flénet et al., 2004; Hartkamp et al., 2002), we chose a sequential optimisation procedure that is fully documented and is easily replicable




**Fig. 5.** Comparison of observed and *STICS* simulated grain yield, grain N, soil water content, and soil mineral N content for calibration and evaluation dataset. rMBE = relative mean bias error, EF = model efficiency, rRMSE = relative Root Mean Square Error. The black line is the 1:1 line. The dotted line represent the regression of simulated against observed values (plotted when significant at 5% risk level). White symbols correspond to the end of growing season.

for other crops and cultivars with the *STICS* model. Breaking down the optimisation into several stages and using it at the module-scale rather than the scale of the entire model allow to avoid unrealistic simulations where errors in parameter values would offset each other (Bechini et al., 2006; Grant, 2001). Taking into account within-season dynamic measurements to successively calibrate the different model components allows to estimate few parameters simultaneously and therefore to

reduce numerical problems (Guillaume et al., 2011; Wallach et al., 2011). The lower and upper boundaries fixed for each parameter during the optimisation ensures that the obtained values correspond to reasonable physiological values thus limiting compensations due to model structure weaknesses. The different optimisation stages and their order following crop development (i.e. phenology, LAI, biomass production and then partitioning) ensure that the parameterisation is relevant




**Fig. 6.** Correlation (a) between grain yield and average water supply/demand ratio during grain onset (over the six days before beginning of grain filling) and (b) total N<sub>2</sub> fixed at harvest and average water deficit factor for N<sub>2</sub> fixation during grain filling (over the first 20 days after beginning of grain filling). Calibration and evaluation datasets were pooled together.




**Fig. 7.** Example of contrasted simulated temporal dynamic (black line) of water supply/demand ratio, above ground biomass and number of grain per square meter in Auch in 2007 (a,d,g), Auzeville in 2010 (b,e,h) and Auzeville in 2007 (c,f,i) in the calibration dataset. White squares are the observations. Vertical dotted lines delimit the periods during which the correlation between average water supply/demand ratio and grain yield is maximal.

according to plant functioning. The order chosen is consistent with previously published calibrations using *STICS* (Corre-Hellou et al., 2009; Flénet et al., 2004) and *DSSAT* (Hartkamp et al., 2002). Finally this method led to a satisfactory match of observed and simulated values, with ranges for statistical indicators (EF, rMBE and rRMSE) consistent with those obtained for the adaption of *STICS* to wheat and maize (Brisson et al., 2002), linseed (Flénet et al., 2004) and spring pea (Corre-Hellou et al., 2009). This approach can be valuable to calibrate new annual crops and other grain legumes (e.g. winter pea, lentil and chick pea) for the *STICS* model, and probably other generic soil-crop models that have a common structure for all species, e.g. *CROPSYST* model (Stöckle et al., 2003).

#### 4.2. A robust and credible agronomic diagnosis

Agronomic diagnosis is crucial for the identification of constraints affecting crop growth and to help the identification of sound agronomic management. The diagnosis needs to be robust, i.e. reliable under

different sets of experimental conditions (Sinclair and Seligman, 2000), and credible, i.e. offering trustworthy causal explanations (Cash et al., 2002). Robustness can be ensured by an explicit reference to physical processes in the construction of mathematical relationships so that the model accurately captures the underlying processes driving the system (Bellocchi et al., 2010). Lack of robustness can lead to a model being “right for the wrong reasons” (Bellocchi et al., 2010). In this section, we want to highlight that we averted these pitfalls, ensuring that our diagnosis was robust and credible.

Our study adds to the body of literature showing that in a temperate climate with rain fed conditions and appropriate control of pest pressure, drought is the prevailing stress and the main driver of yield variability (Karrou and Oweis, 2012; Mafakheri et al., 2010). The explanations we found to explain yield variability (water stress during grain onset) and N<sub>2</sub> fixation variability (water deficit factor during grain filling) are supported by principles of crop physiology derived from experiments and knowledge on response of legume grain yield and N<sub>2</sub> fixation to water stress. With regard to grain yield, water deficit




**Fig. 8.** Example of contrasted simulated temporal dynamic (black line) of water deficit factor for N<sub>2</sub> fixation and N<sub>2</sub> fixed in Auzeville in 2014 (a, c) and 2015 (b, d) in the evaluation dataset. **White** squares are the observations. Vertical dotted lines delimit the periods during which the correlation between average Water deficit factor and N<sub>2</sub> fixation is maximal.

during grain onset reduces grain legume biomass accumulation (decrease of photosynthesis and reduction of leaf area) and assimilate availability that in turn reduce final seed number (Guilioni et al., 2003; Jiang and Egli, 1995; Lake and Sadras, 2016). With regard to N<sub>2</sub> fixation, water stress directly reduce faba bean nodules activity (Sprent, 1972) and sensitivity of N<sub>2</sub> fixation to drought is often maximum during grain filling (Chalk et al., 2010; Mastrodomenico et al., 2013). These two stress responses are implemented in STICS. Firstly, final seed number depends on plant growth rate during the grain onset period (see section 2.1.1 and Table S1). Thus, the magnitude of water stress during grain set and its impact on plant growth rate determines the extent to which the plant can reach the maximum grain number (*nbgrmax*, see Table 4 and Fig. 2). Secondly, in the model, water deficit reduces the potential N<sub>2</sub> fixation, i.e. nodulation and nitrogen fixation are partially or totally inhibited according to nitrate concentration in the topsoil and when soil moisture in the topsoil drops below permanent wilting point (see Section 2.1.2 and Table S1). This careful check of the behaviour of different model functional modules (Figs. 7 and 8) strongly suggests that the STICS model accurately captured the underlying processes driving grain yield formation, N<sub>2</sub> fixation and water stress. We can therefore be confident in the diagnosis drawn from our site-year-management units with regard to water stress. Thus, the generic STICS processes describing yield formation and water stress, when calibrated for faba bean with the proper options activated, have proven useful in explaining variability in yield for this particular legume crop, without the need to build new processes into the generic model.

The good fit obtained with the evaluation dataset indicates that the current set of parameters could be reliably used in other situations different from the ones in the calibration. This however needs to be confirmed with other studies, e.g. for situations where soil mineral N levels would be higher than in our study (on average 47 kg ha<sup>-1</sup> at sowing over the whole profile), which could affect N<sub>2</sub> fixation. We verified by virtual experiment - corresponding to a local sensitivity analysis - that STICS was able to simulate a strong reduction in N<sub>2</sub> fixation when soil nitrate content was higher in the upper layers.

Eventually, by considering both a water deficit factor for N<sub>2</sub> fixation and a water supply/demand ratio (acting on crop biomass), the model allowed unravelling interesting situations where similar faba bean biomass masked differences in the amounts of fixed N<sub>2</sub>. For example, crop biomass and water/supply demand ratio were similarly low in Auzeville in 2011 (3 t ha<sup>-1</sup>) and 2015 (4.1 t ha<sup>-1</sup>) due to similar water stress on biomass (water supply-demand ratio = 0.3). However, the

amount of N<sub>2</sub> fixed was doubled (88 kg ha<sup>-1</sup>) in 2015 compared with 2011 (46 kg ha<sup>-1</sup>). The model indicated a stronger water deficit for N<sub>2</sub> fixation in 2011 (water deficit factor = 0.12) compared with 2015 (water deficit factor = 0.32). This understanding of nonlinearities is crucial for the assessment of innovative cropping system.

Pests and diseases can also reduce faba bean grain yield. Coupling STICS with an epidemiological model (Donatelli et al., 2017) would be relevant to refine the diagnosis in situations where damages on leaves (e.g. rust, *Uromyces viciae-fabae*, or chocolate spot, *Botrytis fabae*) and nodules (e.g. sitonia beetles, *Sitonia lineatus*) can occur.

#### 4.3. New opportunities to design and assess resilient cropping systems

Though the effects of climate change on cereals in Europe have deserved a great attention (e.g. Bregaglio et al., 2017; Dettori et al., 2017), studies on effects of climate change on legume productivity remain scarce. Climate change projections for the temperate and mediterranean regions of Europe point to a decrease in precipitations during spring and summer (Gao and Giorgi, 2008; Giannakopoulos et al., 2009). Water stress will therefore become even more acute with future climate. Our model calibration offers a good opportunity to assess the impact of increased drought-induced stress on faba bean productivity and therefore the potential contribution of legume to sustainable cropping systems in the future.

Dynamic models have often been criticised for putting too much emphasis on single crop assessment on a year-to-year basis (Reckling et al., 2016). Taking into account pre-crop and rotational effects (e.g. nutrient carry-over and alteration of soil water content) will be crucial to accurately assess the contribution of faba bean and grain legumes in building more diversified and resilient cropping systems. Our robust crop calibration and STICS ability to simulate residues mineralisation (Justes et al., 2009), nitrate leaching and water drainage (Plaza-Bonilla et al., 2015), would allow deriving probabilities to achieve a certain level of yield and N provision to the subsequent crop when inserting legumes in cropping systems.

STICS can also simulate intercropping (Brisson et al., 2008). Cereal-legume intercropping (e.g. wheat and faba bean) is a promising integration of legumes in cropping systems in particular in low input cropping systems (Bedoussac et al., 2015). Many processes (e.g. soil N and water sharing) are simulated similarly in sole crop and intercrops (Corre-Hellou et al., 2009), our calibration would therefore allow exploring intercropping performance for a diversity of soils and rainfall

patterns. Nevertheless, some formalisms specific to intercrops (e.g. competition for light) and the corresponding parameters may deserve potential re-calibration before the set of parameters proposed here for sole crops can be used.

## 5. Conclusion

Based on literature and cropping experiments in two contrasting sites of southwestern France, we successfully adapted the *STICS* soil-crop model to faba bean. We developed and validated a method smartly combining the mobilization of literature references, direct field measurements and a transparent and sequential optimisation procedure to derive crop parameters. The obtained set of parameters allowed a satisfactory prediction of plant growth,  $N_2$  fixation, mineral N uptake, grain yield and grain N content. Careful assessment of dynamic variables (plant growth, grain number,  $N_2$  fixation) and the associated stress factors allowed to carry-out a credible and robust diagnosis of grain yield and  $N_2$  fixation variability in the sites of this study. The proposed set of parameter could then be used for studies on (i) yield gap in organic fields affected by biotic and abiotic stresses, (ii) the effect of climate change and more frequent droughts on faba bean, and (iii) the effect of faba bean insertion in crop rotations (as sole crop or intercrop) on N provision to the subsequent crop. Calibration of other grain legumes like chickpea and lentils using the proposed methodology will broaden the capacity of the *STICS* model to simulate innovative cropping systems including grain legumes.

## Acknowledgments

This research was supported by the European Commission (REA) through the LEGATO project (FP7-613551) and the French National Research Agency (ANR) through the LEGITIMES French project (ANR-13-AGRO-0004), the 7<sup>th</sup> PIREN-Seine program and the Climate-CAFE European project (selected by the European FACCE-JPI ERA-NET Plus program). We thank the Centre Régional de Recherche et d'Expérimentation en Agriculture Biologique de Midi-Pyrénées (CREAB-MP) in Auch, France, for making available their faba bean dataset. We are grateful to Loïc Prieur, Didier Raffailac, Michel Labarrère and several trainees who assisted in data collection and Didier Chesneau and Eric Lecloux who performed the extraction and the analysis of soil mineral N. We thank Patrice Lecharpentier for his help with the *Optimistics* software.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.eja.2019.01.001>.

## References

Affholder, F., Scopel, E., Neto, J., Capillon, A., 2003. Diagnosis of the productivity gap using a crop model. Methodology and case study of small-scale maize production in central Brazil. *Agronomie* 23, 305–325.

Amir, J., Sinclair, T.R., 1991. A model of the temperature and solar-radiation effects on spring wheat growth and yield. *Field Crops Res.* 28, 47–58. [https://doi.org/10.1016/0378-4290\(91\)90073-5](https://doi.org/10.1016/0378-4290(91)90073-5).

Anothai, J., Patanothai, A., Jogloy, S., Pannangpetch, K., Boote, K.J., Hoogenboom, G., 2008. A sequential approach for determining the cultivar coefficients of peanut lines using end-of-season data of crop performance trials. *Field Crops Res.* 108, 169–178. <https://doi.org/10.1016/j.fcr.2008.04.012>.

Beaudoin, N., Launay, M., Saubou, E., Ponsardin, G., Mary, B., 2008. Evaluation of the soil crop model STICS over 8 years against the “on farm” database of Bruyères catchment. *Eur. J. Agron.* 29, 46–57. <https://doi.org/10.1016/j.eja.2008.03.001>.

Bécel, C., Munier-Jolain, N.M., Nicolardot, B., 2015. Assessing nitrate leaching in cropping systems based on integrated weed management using the STICS soil-crop model. *Eur. J. Agron.* 62, 46–54. <https://doi.org/10.1016/j.eja.2014.09.005>.

Bechini, L., Bocchi, S., Maggiore, T., Confalonieri, R., 2006. Parameterization of a crop growth and development simulation model at sub-model components level. An example for winter wheat (*Triticum aestivum* L.). *Environ. Model. Softw.* 21, 1042–1054. <https://doi.org/10.1016/j.envsoft.2005.05.006>.

Bedoussac, L., Justes, E., 2010. The efficiency of a durum wheat-winter pea intercrop to improve yield and wheat grain protein concentration depends on N availability during early growth. *Plant Soil* 330, 19–35. <https://doi.org/10.1007/s11104-009-0082-2>.

Bedoussac, L., Journe, E.-P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E.S., Prieur, L., Justes, E., 2015. Ecological principles underlying the increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agron. Sustain. Dev.* 35, 911–935.

Bellocchi, G., Rivington, M., Donatelli, M., Matthews, K., 2010. Validation of biophysical models: issues and methodologies. A review. *Agron. Sustain. Dev.* 30, 109–130. <https://doi.org/10.1051/agro/2009001>.

Bergez, J.E., Raynal, H., Launay, M., Beaudoin, N., Casellas, E., Caubel, J., Chabrier, P., Coucheney, E., Dury, J., García de Cortazar-Atauri, I., Justes, E., Mary, B., Ripoche, D., Ruget, F., 2014. Evolution of the STICS crop model to tackle new environmental issues: new formalisms and integration in the modelling and simulation platform RECORD. *Environ. Model. Softw.* 62, 370–384. <https://doi.org/10.1016/j.envsoft.2014.07.010>.

Bishop, J., Potts, S.G., Jones, H.E., 2016. Susceptibility of faba bean (*Vicia faba* L.) to heat stress during floral development and anthesis. *J. Agron. Crop Sci.* 202, 508–517. <https://doi.org/10.1111/jac.12172>.

Boote, K.J., Mínguez, M.I., Sau, F., 2002. Adapting the CROPGRO legume model to simulate growth of faba bean. *Agron. J.* 94, 743–756.

Bregaglio, S., Hossard, L., Cappelli, G., Resmond, R., Bocchi, S., Barbier, J.-M., Ruget, F., Delmotte, S., 2017. Identifying trends and associated uncertainties in potential rice production under climate change in Mediterranean areas. *Agric. For. Meteorol.* 237–238, 219–232. <https://doi.org/10.1016/j.agrformet.2017.02.015>.

Brisson, N., Mary, B., Ripoche, D., Jeuffroy, M.H., Ruget, F., Nicoulaud, B., Gate, P., Devienne-Barret, F., Antonioletti, R., Durr, C., et al., 1998. STICS: a generic model for the simulation of crops and their water and nitrogen balances. I. Theory and parameterization applied to wheat and corn. *Agronomie* 18, 311–346.

Brisson, N., Ruget, F., Gate, P., Lorgeou, J., Nicoulaud, B., Tayot, X., Plenet, D., Jeuffroy, M.-H., Bouthier, A., Ripoche, D., Mary, B., Justes, E., 2002. STICS: a generic model for simulating crops and their water and nitrogen balances. II. Model validation for wheat and maize. *Agronomie* 22, 69–92. <https://doi.org/10.1051/agro:2001005>.

Brisson, N., Launay, M., Mary, B., Beaudoin, N., 2009. Conceptual Basis, Formalisations and Parameterization of the Stics Crop Model. Editions Quae.

Bruelle, G., Affholder, F., Abrell, T., Ripoche, A., Dusserre, J., Naudin, K., Titttonell, P., Rabeharisoa, L., Scopel, E., 2017. Can conservation agriculture improve crop water availability in an erratic tropical climate producing water stress? A simple model applied to upland rice in Madagascar. *Agric. Water Manag.* 192, 281–293. <https://doi.org/10.1016/j.agwat.2017.07.020>.

Buis, S., Wallach, D., Guillaume, S., Varella, H., Lecharpentier, P., Launay, M., Guerif, M., Berge, J.-E., Justes, E., 2011. The STICS crop model and associated software for analysis, parameterization, and evaluation. Methods of Introducing System Models into Agricultural Research - Advances in Agricultural Systems Modeling 2. L.R. Ahuja Liwang Ma, pp. 395–426.

Casa, R., Varella, H., Buis, S., Guérif, M., De Solan, B., Baret, F., 2012. Forcing a wheat crop model with LAI data to access agronomic variables: evaluation of the impact of model and LAI uncertainties and comparison with an empirical approach. *Eur. J. Agron.* 37, 1–10. <https://doi.org/10.1016/j.eja.2011.09.004>.

Cash, D., Clark, W.C., Alcock, F., Dickson, N.M., Eckley, N., Jäger, J., 2002. Salience, Credibility, Legitimacy and Boundaries: Linking Research, Assessment and Decision Making (SSRN Scholarly Paper No. ID 372280). Social Science Research Network, Rochester, NY.

Chalk, P.M., Alves, B.J.R., Boddey, R.M., Urquiga, S., 2010. Integrated effects of abiotic stresses on inoculant performance, legume growth and symbiotic dependence estimated by 15N dilution. *Plant Soil* 328, 1–16. <https://doi.org/10.1007/s11104-009-0187-7>.

Colomb, B., Jouany, C., Prieur, L., 2013. Des bilans de phosphore majoritairement négatifs pour les systèmes de grandes cultures biologiques sans élevage en Midi-Pyrénées. Quels impacts sur le phosphore biodisponible des sols et l'état de nutrition des cultures? *Innov. Agron.* 32, 73–82.

Confalonieri, R., Bechini, L., 2004. A preliminary evaluation of the simulation model CropSyst for alfalfa. *Eur. J. Agron.* 21, 223–237. <https://doi.org/10.1016/j.eja.2003.08.003>.

Constantin, J., Dürr, C., Tribouillois, H., Justes, E., 2015a. Catch crop emergence success depends on weather and soil seedbed conditions in interaction with sowing date: a simulation study using the SIMPLE emergence model. *Field Crops Res.* 176, 22–33. <https://doi.org/10.1016/j.fcr.2015.02.017>.

Constantin, J., Le Bas, C., Justes, E., 2015b. Large-scale assessment of optimal emergence and destruction dates for cover crops to reduce nitrate leaching in temperate conditions using the STICS soil-crop model. *Eur. J. Agron.* 69, 75–87. <https://doi.org/10.1016/j.eja.2015.06.002>.

Corre-Hellou, G., Faure, M., Launay, M., Brisson, N., Crozat, Y., 2009. Adaptation of the STICS intercrop model to simulate crop growth and N accumulation in pea-barley intercrops. *Field Crops Res.* 113, 72–81. <https://doi.org/10.1016/j.fcr.2009.04.007>.

Coucheney, E., Buis, S., Launay, M., Constantin, J., Mary, B., García de Cortazar-Atauri, I., Ripoche, D., Beaudoin, N., Ruget, F., Andrianarisoa, K.S., Le Bas, C., Justes, E., Léonard, J., 2015. Accuracy, robustness and behavior of the STICS soil-crop model for plant and nitrogen outputs: evaluation over a wide range of agro-environmental conditions in France. *Environ. Model. Softw.* 64, 177–190. <https://doi.org/10.1016/j.envsoft.2014.11.024>.

Daryanto, S., Wang, L., Jacinthe, P.-A., 2017. Global synthesis of drought effects on cereal, legume, tuber and root crops production: a review. *Agric. Water Manag.* 179, 18–33. <https://doi.org/10.1016/j.agwat.2016.04.022>.

Dettoni, M., Cesaraccio, C., Duce, P., 2017. Simulation of climate change impacts on

- production and phenology of durum wheat in Mediterranean environments using CERES-Wheat model. *Field Crops Res.* 206, 43–53. <https://doi.org/10.1016/j.fcr.2017.02.013>.
- Di Paola, A., Valentini, R., Santini, M., 2016. An overview of available crop growth and yield models for studies and assessments in agriculture. *J. Sci. Food Agric.* 96, 709–714. <https://doi.org/10.1002/jsfa.7359>.
- Donatelli, M., Magarey, R.D., Bregaglio, S., Willocquet, L., Whish, J.P.M., Savary, S., 2017. Modelling the impacts of pests and diseases on agricultural systems. *Agric. Syst.* 155, 213–224. <https://doi.org/10.1016/j.agry.2017.01.019>.
- Ellis, R.H., Summerfield, R.J., Roberts, E.H., 1988. Effects of Temperature, Photoperiod and Seed Vernalization on Flowering in Faba Bean *Vicia faba*. *Ann. Bot.* 61, 17–27. <https://doi.org/10.1093/oxfordjournals.aob.a087524>.
- Flénet, F., Villon, P., Ruget, F., 2004. Methodology of adaptation of the STICS model to a new crop: spring linseed (*Linum usitatissimum*, L.). *Agronomie* 24, 367–381. <https://doi.org/10.1051/agro:2004032>.
- Flores, F., Nadal, S., Solis, I., Winkler, J., Sass, O., Stoddard, F.L., Link, W., Raffiot, B., Muel, F., Rubiales, D., 2012. Faba bean adaptation to autumn sowing under European climates. *Agron. Sustain. Dev.* 32, 727–734. <https://doi.org/10.1007/s13593-012-0082-0>.
- Gao, X., Giorgi, F., 2008. Increased aridity in the Mediterranean region under greenhouse gas forcing estimated from high resolution simulations with a regional climate model. *Glob. Planet. Change* 62, 195–209. <https://doi.org/10.1016/j.gloplacha.2008.02.002>.
- Giannakopoulos, C., Le Sager, P., Bindi, M., Moriondo, M., Kostopoulou, E., Goodess, C.M., 2009. Climatic changes and associated impacts in the Mediterranean resulting from a 2 °C global warming. *Glob. Planet. Change* 68, 209–224. <https://doi.org/10.1016/j.gloplacha.2009.06.001>.
- Grant, F.R., 2001. Modeling transformations of soil organic carbon and nitrogen at different scales of complexity. In: Ma, L., Hansen, S., Shaffer, M.J. (Eds.), *Modeling Carbon and Nitrogen Dynamics for Soil Management*. Lewis Publishers.
- Guilioni, L., Wery, J., Lecoer, J., 2003. High temperature and water deficit may reduce seed number in field pea purely by decreasing plant growth rate. *Funct. Plant Biol.* 30, 1151–1164. <https://doi.org/10.1071/fp03105>.
- Guillaume, S., Bergez, J.-E., Wallach, D., Justes, E., 2011. Methodological comparison of calibration procedures for durum wheat parameters in the STICS model. *Eur. J. Agron.* 35, 115–126. <https://doi.org/10.1016/j.eja.2011.05.003>.
- Hartkamp, A.D., Hoogenboom, G., White, J.W., 2002. Adaptation of the CROPGRO growth model to velvet bean (*Mucuna pruriens*): I. Model development. *Field Crops Res.* 78, 9–25. [https://doi.org/10.1016/S0378-4290\(02\)00091-6](https://doi.org/10.1016/S0378-4290(02)00091-6).
- Hayman, P.T., Whitbread, A.M., Gobbett, D.L., 2010. The impact of El Niño Southern Oscillation on seasonal drought in the southern Australian grainbelt. *Crop Pasture Sci.* 61, 528–539. <https://doi.org/10.1071/CP09221>.
- He, D., Wang, E., Wang, J., Robertson, M.J., 2017. Data requirement for effective calibration of process-based crop models. *Agric. For. Meteorol.* 234–235, 136–148. <https://doi.org/10.1016/j.agrformet.2016.12.015>.
- Jégo, G., Pattey, E., Bourgeois, G., Morrison, M.J., Drury, C.F., Tremblay, N., Tremblay, G., 2010. Calibration and performance evaluation of soybean and spring wheat cultivars using the STICS crop model in Eastern Canada. *Field Crops Res.* 117, 183–196. <https://doi.org/10.1016/j.fcr.2010.03.008>.
- Jensen, E.S., Peoples, M.B., Hauggaard-Nielsen, H., 2010. Faba bean in cropping systems. *Field Crops Res.* 115, 203–216. <https://doi.org/10.1016/j.fcr.2009.10.008>.
- Jiang, H., Egli, D.B., 1995. Soybean seed number and crop growth rate during flowering. *Agron. J.* 87, 264–267.
- Jones, J., Hoogenboom, G., Porter, C., Boote, K., Batchelor, W., Hunt, L., Wilkens, P., Singh, U., Gijssman, A., Ritchie, J., 2003. The DSSAT cropping system model. *Eur. J. Agron.* 18, 235–265. [https://doi.org/10.1016/S1161-0301\(02\)00107-7](https://doi.org/10.1016/S1161-0301(02)00107-7).
- Justes, E., Mary, B., Nicolardot, B., 2009. Quantifying and modelling C and N mineralization kinetics of catch crop residues in soil: parameterization of the residue decomposition module of STICS model for mature and non mature residues. *Plant Soil* 325, 171–185. <https://doi.org/10.1007/s11104-009-9966-4>.
- Kammoun, B., 2014. interactions génotype x environnement x conduite culturale de peuplement bi-spécifique de cultures associées de blé dur et de légumineuses à graines, à des fins de choix variétal et d'optimisation de leurs itinéraires techniques. *École Doctorale Sciences Écologiques, Vétérinaires, Agronomiques et Bioingénieries (Toulouse)*, 154236330.
- Karrou, M., Oweis, T., 2012. Water and land productivities of wheat and food legumes with deficit supplemental irrigation in a Mediterranean environment. *Agric. Water Manag.* 107, 94–103. <https://doi.org/10.1016/j.agwat.2012.01.014>.
- Keating, B., Carberry, P., Hammer, G., Probert, M., Robertson, M., Holzworth, D., Huth, N., Hargreaves, J.N., Meinke, H., Hochman, Z., McLean, G., Verburg, K., Snow, V., Dimes, J., Silburn, M., Wang, E., Brown, S., Bristow, K., Asseng, S., Chapman, S., McCown, R., Freebairn, D., Smith, C., 2003. An overview of APSIM, a model designed for farming systems simulation. *Eur. J. Agron.* 18, 267–288. [https://doi.org/10.1016/S1161-0301\(02\)00108-9](https://doi.org/10.1016/S1161-0301(02)00108-9).
- Khan, H.R., Paull, J.G., Siddique, K.H.M., Stoddard, F.L., 2010. Faba bean breeding for drought-affected environments: A physiological and agronomic perspective. *Field Crops Res.* 115, 279–286. <https://doi.org/10.1016/j.fcr.2009.09.003>.
- Ko, J., Piccinini, G., Guo, W., Steglich, E., 2009. Parameterization of EPIC crop model for simulation of cotton growth in South Texas. *J. Agric. Sci.* 147, 169–178. <https://doi.org/10.1017/S0021859608008356>.
- Lake, L., Sadras, V.O., 2016. Screening chickpea for adaptation to water stress: associations between yield and crop growth rate. *Eur. J. Agron.* 81, 86–91. <https://doi.org/10.1016/j.eja.2016.09.003>.
- Lemaire, G., Gastal, F., 1997. N uptake and distribution in plant canopies. *Diagnosis of the Nitrogen Status in Crops*. Springer, Berlin, Heidelberg, pp. 3–43. [https://doi.org/10.1007/978-3-642-60684-7\\_1](https://doi.org/10.1007/978-3-642-60684-7_1).
- Liu, Y., Wu, L., Baddeley, J.A., Watson, C.A., 2011. Models of biological nitrogen fixation of legumes. A review. *Agron. Sustain. Dev.* 31, 155–172. <https://doi.org/10.1051/agro/2010008>.
- López-Bellido, R.J., López-Bellido, L., Benítez-Vega, J., Muñoz-Romero, V., López-Bellido, F.J., Redondo, R., 2011. Chickpea and faba bean nitrogen fixation in a Mediterranean rainfed Vertisol: Effect of the tillage system. *Eur. J. Agron.* 34, 222–230. <https://doi.org/10.1016/j.eja.2011.01.005>.
- Mafakheri, A., Siosemardeh, A., Bahramnejad, B., Struik, P.C., Sohrabi, Y., 2010. Effect of drought stress on yield, proline and chlorophyll contents in three chickpea cultivars. *Aust. J. Crop Sci.* 4, 580–585.
- Mastrodomenico, A.T., Purcell, L.C., Andy King, C., 2013. The response and recovery of nitrogen fixation activity in soybean to water deficit at different reproductive developmental stages. *Environ. Exp. Bot.* 85, 16–21. <https://doi.org/10.1016/j.envexpbot.2012.07.006>.
- Meier, U., 2001. Growth Stages of Mono- and Dicotyledonous Plants: BBCH Monograph. Federal Biological Research Centre for Agriculture and Forestry.
- Meynard, J.-M., Jeuffroy, M.-H., Le Bail, M., Lefèvre, A., Magrini, M.-B., Michon, C., 2017. Designing coupled innovations for the sustainability transition of agrifood systems. *Agric. Syst.* 157, 330–339. <https://doi.org/10.1016/j.agry.2016.08.002>.
- Parent, B., Tardieu, F., 2012. Temperature responses of developmental processes have not been affected by breeding in different ecological areas for 17 crop species. *New Phytol.* 194, 760–774. <https://doi.org/10.1111/j.1469-8137.2012.04086.x>.
- Patrick, J.W., Stoddard, F.L., 2010. Physiology of flowering and grain filling in faba bean. *Field Crops Res.* 115, 234–242. <https://doi.org/10.1016/j.fcr.2009.06.005>.
- Plaza-Bonilla, D., Nolot, J.-M., Raffallac, D., Justes, E., 2015. Cover crops mitigate nitrate leaching in cropping systems including grain legumes: field evidence and model simulations. *Agric. Ecosyst. Environ.* 212, 1–12. <https://doi.org/10.1016/j.agee.2015.06.014>.
- Plaza-Bonilla, D., Nolot, J.-M., Raffallac, D., Justes, E., 2017. Innovative cropping systems to reduce N inputs and maintain wheat yields by inserting grain legumes and cover crops in southwestern France. *Eur. J. Agron.* 82 (Part B), 331–341. <https://doi.org/10.1016/j.eja.2016.05.010>.
- Prasad, P.V.V., Bheemanahalli, R., Jagadish, S.V.K., 2017. Field crops and the fear of heat stress—opportunities, challenges and future directions. *Field Crops Res.* 200, 114–121. <https://doi.org/10.1016/j.fcr.2016.09.024>.
- Reckling, M., Hecker, J.-M., Bergkvist, G., Watson, C.A., Zander, P., Schläpke, N., Stoddard, F.L., Eory, V., Topp, C.F.E., Maire, J., Bachinger, J., 2016. A cropping system assessment framework—Evaluating effects of introducing legumes into crop rotations. *Eur. J. Agron.* 76, 186–197. <https://doi.org/10.1016/j.eja.2015.11.005>.
- Rubiales, D., Fondevilla, S., Chen, W., Gentzbitel, L., Higgins, T.J.V., Castillejo, M.A., Singh, K.B., Rispail, N., 2015. Achievements and challenges in legume breeding for pest and disease resistance. *Crit. Rev. Plant Sci.* 34, 195–236. <https://doi.org/10.1080/07352689.2014.898445>.
- Ruget, F., Brisson, N., Delécolle, R., Faivre, R., 2002. Sensitivity analysis of a crop simulation model, STICS, in order to choose the main parameters to be estimated. *Agronomie* 22, 133–158. <https://doi.org/10.1051/agro:2002009>.
- Saxton, K.E., Rawls, W.J., 2006. Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Sci. Soc. Am. J.* 70, 1569–1578. <https://doi.org/10.2136/sssaj2005.0117>.
- Sinclair, T.R., Seligman, N., 2000. Criteria for publishing papers on crop modeling. *Field Crops Res.* 68, 165–172. [https://doi.org/10.1016/S0378-4290\(00\)00105-2](https://doi.org/10.1016/S0378-4290(00)00105-2).
- Sprent, J.I., 1972. The effects of water stress on nitrogen-fixing root nodules. *New Phytol.* 71, 603–611. <https://doi.org/10.1111/j.1469-8137.1972.tb01270.x>.
- Stoate, C., Baldi, A., Beja, P., Boatman, N.D., Herzog, I., van Doorn, A., de Snoo, G.R., Rakosy, L., Ramwell, C., 2009. Ecological impacts of early 21st century agricultural change in Europe – a review. *J. Environ. Manage.* 91, 22–46. <https://doi.org/10.1016/j.jenvman.2009.07.005>.
- Stöckle, C.O., Donatelli, M., Nelson, R., 2003. CropSyst, a cropping systems simulation model. *Eur. J. Agron.* 18, 289–307. [https://doi.org/10.1016/S1161-0301\(02\)00109-0](https://doi.org/10.1016/S1161-0301(02)00109-0).
- Stoddard, F.L., Nicholas, A.H., Rubiales, D., Thomas, J., Villegas-Fernández, A.M., 2010. Integrated pest management in faba bean. *Field Crops Res.* 115, 308–318. <https://doi.org/10.1016/j.fcr.2009.07.002>.
- Tribouillois, H., Dürr, C., Demilly, D., Wagner, M.-H., Justes, E., 2016. Determination of germination response to temperature and water potential for a wide range of cover crop species and related functional groups. *PLoS One* 11. <https://doi.org/10.1371/journal.pone.0161185>.
- Unkovich, M., 2008. Measuring Plant-associated Nitrogen Fixation in Agricultural Systems. Australian Centre for International Agricultural Research, Canberra.
- Wallach, D., Buis, S., Lecharpentier, P., Bourges, J., Clastre, P., Launay, M., Bergez, J.-E., Guerif, M., Soudais, J., Justes, E., 2011. A package of parameter estimation methods and implementation for the STICS crop-soil model. *Environ. Model. Softw.* 26, 386–394. <https://doi.org/10.1016/j.envsoft.2010.09.004>.
- Watson, C.A., Reckling, M., Preissel, S., Bachinger, J., Bergkvist, G., Kuhlman, T., Lindström, K., Nemecek, T., Topp, C.F.E., Vanhatalo, A., Zander, P., Murphy-Bokern, D., Stoddard, F.L., 2017. Grain legume production and use in European agricultural systems. *Adv. Agron.* 144, 235–303. <https://doi.org/10.1016/bs.agron.2017.03.003>.